

AD-A173 019


A PSEUDO-ARCLENGTH CONTINUATION METHOD FOR NONLINEAR
EIGENVALUE PROBLEMS(U) ARIZONA STATE UNIV TEMPE DEPT OF
MATHEMATICS H D MITTELHANN OCT 85 TR-88 AFOSR-84-0315


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A173 019

A Pseudo-Arclength Continuation Method
for Nonlinear Eigenvalue Problems

by

H. D. Mittelmann

Technical Report No. 88
October 1985

Arizona State University
Tempe, Arizona 85287

DEPARTMENT OF MATHEMATICS

Approved for public release;
distribution unlimited.

DTIC
S ELECTE D
OCT 08 1986
D



AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for public release IAW AFR 193-12.
Distribution is unlimited.

WILLIAM J. KEEPER
Chief, Technical Information Division

ARIZONA STATE UNIVERSITY

DTIC FILE COPY

86 10 8 146

1

A Pseudo-Arclength Continuation Method
for Nonlinear Eigenvalue Problems

by

H. D. Mittelmann

Technical Report No. 88
October 1985

Arizona State University
Tempe, Arizona 85287

DTIC
ELECTE
OCT 08 1986
S D

Arizona State University is an affirmative action/equal employment opportunity institution and does not discriminate on the basis of sex, race, ethnic origin, creed, color, age, handicapped or veteran status in its program or employment.

This document is approved
for public release and
distribution is unlimited.

A PSEUDO-ARCLENGTH CONTINUATION METHOD
FOR NONLINEAR EIGENVALUE PROBLEMS

Hans D. Mittelmann[†]

Abstract. A variant of the classical pseudo-arclength continuation method is proposed. Basically, the method can be viewed as pseudo-arclength continuation in (r, λ) space where r is a functional of the solution. Another difference is a three-parameter predictor instead of the standard Euler step. This predictor, as well as the Newton corrector iteration, are justified and some numerical results for reaction-diffusion equations are presented. The method provides a simple algebraic check for symmetry-breaking bifurcation, the most common type of secondary bifurcation in physical examples.

Keywords: parameter-dependent boundary value problems, continuation algorithm; singular points; symmetry-breaking bifurcation; reaction-diffusion equations.

[†] Department of Mathematics, Arizona State University,
Tempe, Arizona 85287

This work was supported by the Air Force Office of
Scientific Research under Grant AFOSR-84-0315



<input checked="" type="checkbox"/>	
<input type="checkbox"/>	
<input type="checkbox"/>	
y Codes	
Dist	and/or Special
A-1	

1. Introduction

Boundary value problems for partial differential equations that describe phenomena in many fields of application are usually nonlinear and depend on one or several parameters of the underlying system. These parameters enter in various ways. The coefficients or right-hand sides of the differential equations may depend on them, the boundary conditions or, for example, the domain. With respect to these parameters, in general, bifurcation and non-unicity effects occur. For a list of different problems from the applications as well as a first introduction to analytic and numerical aspects we refer to [8]. A survey of the state of the art may be obtained from [7].

Frequently, the only way to explore the solution manifolds of these problems is to continue along the solution branches, in the case of multi-parameter dependence, with respect to one of the parameters for fixed values of the others. This approach yields a quantitatively satisfactory approximation if, for example, suitable discretizations are used, while it may be difficult to get a full qualitative impression in this way. Other analytic techniques yielding the latter but not also a quantitatively useful result, may well supplement the numerical methods.

For the continuation along solution branches, different methods have been proposed and applied. We restrict considerations here to the methods introducing an arclength-like additional parameter. One of the earlier and frequently referenced papers is [6]. To be more specific, let the

parameter-dependent nonlinear problem be given by

$$(1.1) \quad G(u, \alpha) = 0$$

where u is an element of a Banach space X and G maps $X \times \mathbb{R}^p$ into X . We denote by λ the component of α that is presently not fixed and, for simplicity, suppress the others. The pseudo-arclength method of [6] then introduces an additional parameter σ and augments (1.1) by a normalizing condition

$$(1.2) \quad \begin{aligned} G(u, \lambda) &= 0, \\ N(u(\sigma), \lambda(\sigma), \sigma) &= 0. \end{aligned}$$

Thus, u as well as λ are locally parametrized by σ . In the following section, we will introduce a slight modification of this approach. In applications, one is frequently interested in a functional of u , for example, a norm, the value of the definite integral of u over some region, etc. This quantity may represent a physically important parameter just as the components of α . For graphical representations also, usually such a functional is depicted versus one or two of the α 's. Using a functional of u , therefore, is not equivalent to introducing another artificial parameter.

After introducing the basic continuation method, we discuss a new predictor to be used in conjunction with it in Section 3. An analysis of the corrector iteration is given in the following section. and in the last section, some numerical results are presented.

2. The Continuation Method

For the problem

$$(2.1) \quad G(u, \lambda) = 0, \quad G: X \times R \rightarrow X$$

let s denote the arclength along a solution arc $(u(s), \lambda(s))$. Under sufficient smoothness and regularity assumptions this latter dependence is differentiable and thus $(G_u^0 = G_u(u_0, \lambda_0))$

$$(2.2a) \quad G_u^0 \dot{u}_0 + G_\lambda^0 \dot{\lambda}_0 = 0,$$

$$(2.2b) \quad \|\dot{u}_0\|^2 + \dot{\lambda}_0^2 = 1$$

in a solution $(u_0, \lambda_0) = (u(s_0), \lambda(s_0))$ of (2.1). Here, subscripts stand for partial derivatives and a dot for differentiation with respect to s .

In [6] it was proposed to augment (2.1) by the following normalizing condition:

$$(2.3) \quad N_\sigma(u, \lambda, \sigma) \equiv \theta \dot{u}_0(u - u_0) + (2 - \theta) \dot{\lambda}_0(\lambda - \lambda_0) - \delta\sigma = 0$$

which approximates (2.2b). σ is thus called the pseudo-arclength parameter. $\delta\sigma$ is the steplength in σ . θ satisfying $0 < \theta < 2$ was meant to be a weighting parameter.

Let $r(u)$ denote a functional of u . For simplicity, let

$$(2.4) \quad r(u) = \|u\|$$

where $\|\cdot\|$ is the norm of u which we assume to be differentiable. Other functionals may be chosen, as for example, in VLSI problems ([2]), but in the following we restrict considerations to (2.4). Quite frequently, this parameter is used in a graphical representation of the solution manifold and also has a physical significance.

The following augmenting equation leads to a pseudo-arclength method in (r, λ) -space.

$$(2.5) \quad N_r(u, \lambda, \sigma) \equiv \theta \dot{r}_0(r - r_0) + (2 - \theta) \dot{\lambda}_0(\lambda - \lambda_0) - \delta\sigma = 0,$$

where $0 \leq \theta \leq 2$, $r_0 = \|u_0\|$, $\dot{r}_0 = \left(\frac{d}{d\sigma} \|u\|\right)_\sigma = \sigma_0$.

While the classical pseudo-arclength method with Euler predictor using (2.3) is in general optimal in a local sense among continuation procedures utilizing only first derivatives and depending only on one parameter, in this case the step-length along the tangent, (2.5) will be combined with a more expensive predictor to yield in many examples a better global continuation behaviour. This will be demonstrated in 5.

Let us assume from now on that X is a Hilbert space, sufficient regularity is given, and $\|\cdot\| = (\cdot, \cdot)^{1/2}$, then

$$(2.6) \quad \dot{r}_0 = (u_0, \dot{u}_0) / r_0.$$

From (2.5) we see that the three values 0, 1, 2 for θ are special in the sense that for $\theta = 0$ (2) the augmenting equation characterizes points with a fixed $\lambda(r)$ -value, while for $\theta = 1$

the points lie on a hyperplane orthogonal to the solution arc in (r_0, λ_0) .

This suggests to continue to target values in the two physical parameters r and λ . Such a strategy seems appropriate as long as starting from a point on the solution curve another point corresponding to a suitably picked target value can be computed in a few number of (corrector) iterations. This will, of course, depend on the starting guess and thus on the method used for predicting this point. Only the combination of (2.5) with the predictor presented in the next section makes this continuation the effective and powerful algorithm it has proven to be at least for a series of problems from the applications.

3. The Predictor

In continuing from a point (u_0, λ_0) on a solution branch most continuation methods first generate a starting guess (u_p, λ_p) and then iteratively determine a new solution point. The pseudo-arclength method in its standard version uses a first order Euler predictor

$$(3.1) \quad \begin{pmatrix} u_p \\ \lambda_p \end{pmatrix} = \begin{pmatrix} u_0 + \delta \sigma \dot{u}_0 \\ \lambda_0 + \delta \sigma \dot{\lambda}_0 \end{pmatrix} .$$

Subsequently, the augmented system (2.1), (2.3) is iteratively solved ('corrector'). The computed solution for $\theta = 1$ is thus located in the intersection of the solution manifold and a hyperplane perpendicular to the tangent vector $(\dot{u}_0, \dot{\lambda}_0)^T$ at the

previous point and a distance $\delta\sigma$ from this point. It is not immediately apparent how the parameter θ would have to be chosen to improve the quality of the predicted point.

It is clear that with respect to the information used, namely \dot{u}_0 and $\dot{\lambda}_0$, the Euler predictor step is locally, i.e., for $\delta\sigma \rightarrow 0$, optimal, but that, for example, in highly curved parts of the solution arc, only a rather small $\delta\sigma$ will lead to convergence of the corrector iteration.

In order to improve the quality of the predictor either a higher order method making use of higher derivatives of G may be used or in some other way, it has to be assured that the predicted point better approximates a solution of the nonlinear eigenvalue problem. The first approach is suitable in particular if the nonlinearities have a simple form as, for example, low order polynomials. Here, we propose to make use of only the first order information also used in the Euler predictor. Instead of a single parameter, three are used in

$$(3.2) \quad \begin{pmatrix} u_p \\ \lambda_p \end{pmatrix} = \begin{pmatrix} (1+\gamma)u_0 + \beta\dot{u}_0 \\ \lambda_0 + \alpha\dot{\lambda}_0 \end{pmatrix}.$$

In case $\alpha \neq \beta$, $\gamma = 0$ the predicted point is in a non-tangential direction from (u_0, λ_0) . This is slightly more general than (3.1) but, of course, treats λ as a special variable as opposed to methods that, for a finite-dimensional problem, give λ no preferential treatment ([12]). The justification for (3.2) can only be that λ , in fact, represents an important physical parameter.

More explanation is needed for the coefficient of u_0 in (3.2). Before this will be done, we complete the definition of the predictor. A system of three equations is used to determine the parameters α , β and γ .

$$\begin{aligned} N(u_p, \lambda_p, \sigma) &= 0, \\ (3.3) \quad (u_0, G(u_p, \lambda_p)) &= 0, \\ (\dot{u}_0, G(u_p, \lambda_p)) &= 0. \end{aligned}$$

In addition to the normalizing equation, two orthogonality relations with respect to the scalar product on X are imposed. The basis for these equations is twofold. First, it assures that the predicted point is a weak solution of (2.1) with respect to the subspace spanned by u_0 and \dot{u}_0 . On the other hand, (3.2) shows that

$$(3.4) \quad (u_p, G(u_p, \lambda_p)) = 0$$

holds, which defines a generalized Rayleigh-quotient.

The system (3.3) in general need not have a solution (α, β, γ) or such a solution is not necessarily unique. If, for example, u_0 and \dot{u}_0 are parallel, then the rank of the Jacobian of (3.3) is at most two. Therefore, a least squares solution of (3.3) is to be computed. In order to be able to analyze several important features of the continuation method, we have

to more closely consider the case of bifurcation points. Let us introduce the notation

$$(3.5) \quad F(y) = \begin{pmatrix} G(u, \lambda) \\ N(u, \lambda, \sigma) \end{pmatrix}, \quad y = (u, \lambda)$$

and use subscripts for partial derivatives. (u_0, λ_0) is a simple pitchfork (symmetric) bifurcation point of (2.1) if the following conditions hold

$$(3.6) \quad \begin{aligned} N(G_u^0) &= \text{span} \{ \phi_0 \}, \quad \|\phi_0\| = 1, \\ R(G_u^0) &\text{ is closed and } \text{codim } R(G_u^0) = 1, \\ \phi_0^* G_\lambda^0 &= 0, \quad \phi_0^* G_{uu}^0 \phi_0 \phi_0 = 0, \\ b &= \phi_0^* (G_{u\lambda}^0 \phi_0 + G_{uu}^0 v_0 \phi_0) \neq 0 \end{aligned}$$

where v_0 is the unique solution of

$$(3.7) \quad G_u^0 v_0 + G_\lambda^0 = 0, \quad \phi_0^* v_0 = 0.$$

N, R denote nullspace and range, respectively, and

$$N(G_u^{0*}) = \text{span} \{ \phi_0^* \}, \quad \phi_0^* \phi_0 = 1.$$

From differentiating (2.1) twice with respect to s it follows that (cf. [4]) the algebraic bifurcation equation

$$(3.8) \quad (c\dot{\lambda}_0 + 2b\xi_1)\dot{\lambda}_0 = 0$$

holds where $c = \phi_0^*(G_{uu}^0 v_0 v_0 + 2G_{u\lambda}^0 v_0 + G_{\lambda\lambda}^0)$ and $\xi_1^2 + 2\dot{\lambda}_0^2 = 1$. The two solution branches passing through (u_0, λ_0) are thus given by

$$(3.9) \quad \begin{aligned} u(s) &= u_0 + (s-s_0)(\dot{\lambda}_0 v_0 + \xi_1 \phi_0) + O((s-s_0)^2), \\ \lambda(s) &= \lambda_0 + (s-s_0) \dot{\lambda}_0 + O((s-s_0)^2). \end{aligned}$$

These pitchfork bifurcation points are not generic in the sense that they are in general not present under perturbations of the problem as, for example, introduced by a discretization. An even further specialized class of bifurcation points, the symmetry-breaking bifurcation points, however, have this property.

Let us assume for simplicity that S is a linear operator on X , $S^2 = I$, $S \neq I$ such that

$$(3.10) \quad G(Su, \lambda) = SG(u, \lambda), \text{ for all } u \in X, \lambda \in \mathbb{R}.$$

Then X may be split as

$$(3.11) \quad X = X_s \oplus X_a,$$

where $X_s = \{u \in X, u = Su\}$ contains the symmetric elements with respect to S and $X_a = \{u \in X, u = -Su\}$, the anti-symmetric elements. The pitchfork bifurcation point (u_0, λ_0) is then symmetry-breaking if

$$(3.12) \quad u_0 \in X_s, \quad v_0 \in X_s, \quad \phi_0 \in X_a.$$

On the branch corresponding to the solution $\dot{\lambda}_0 = 0$ of (3.8) it thus holds that

$$(3.13) \quad \dot{u}_0 = \pm \phi_0 \in X_a$$

while on the other branch $c = 0$ implies $\dot{u}_0 \in X_s$. On the symmetry-breaking branch we have from (2.6) if $S = S^*$

$$(3.14) \quad \dot{\lambda}_0 = 0, \quad \dot{r}_0 = 0$$

and thus, in this case, N_r in (2.5) is not well defined. The restriction to the two-dimensional (r, λ) -subspace introduces additional singularities that are not present in the classical pseudo-arclength method. A natural modification is to introduce as a parameter the norm of the anti-symmetric part u_a of u corresponding to the splitting (3.11). This was done explicitly in [10]. Since on the branch of nonsymmetric solutions $u_a = \dot{u}_0 \in X_a$, asymptotically in the bifurcation point from (3.9), we propose instead in the general case to use $N = N_\sigma$ in (3.3) whenever $|\dot{r}_0|/\|\dot{u}_0\|$ is small.

Let us assume now that starting at (u_0, λ_0) a point (u, λ) on the symmetry-breaking branch with $r = \|u\| < r_0 = \|u_0\|$ is to be computed. For $\gamma = 0$ (3.2), (3.12), (3.13) imply that always $\|u_p\| > r_0$. This is one of the reasons that γ was

introduced in (3.2). Another reason is that in the linear eigenvalue problem, computations along the eigenfunction branches require a renormalization only and are not prone to cancellation if in the predictor step u_p is actually obtained by renormalization, i.e., $\beta = 0$ in (3.2).

Let us summarize the normalizing condition that is proposed in the various cases.

$$\begin{aligned}
 & |\dot{r}_0| < \epsilon \|\dot{u}_0\|: & N = N_\sigma, \theta \neq 0 \\
 (3.15) \quad & |\dot{r}_0| > (1-\epsilon)\|\dot{u}_0\|: & N = N_r, \theta \neq 0, \beta = 0, \\
 & \epsilon \|\dot{u}_0\| \leq \dot{r}_0 \leq (1-\epsilon)\|\dot{u}_0\|: & N = N_r, 0 \leq \theta \leq 2.
 \end{aligned}$$

In the following section, we will address the question of the regularity of the corrector Jacobian in the various cases of (3.15).

4. The Corrector

The point (u_p, λ_p) computed by the predictor described in the previous section is used as usual as starting guess for an iterative solution of the augmented system (3.5). This is done by Newton's method and thus the Jacobian

$$(4.1) \quad F_y = \begin{pmatrix} G_u & G_\lambda \\ N_u & N_\lambda \end{pmatrix}$$

has to be evaluated and inverted. The following lemma is a simplified version of Lemma 2.8 in [6].

Lemma 4.1 Let X be a Banach space and consider the linear operator $\bar{A}: X \times R \rightarrow X \times R$ of the form

$$\bar{A} = \begin{pmatrix} A & B \\ C^* & D \end{pmatrix}$$

where $A: X \rightarrow X$, $B: R \rightarrow X$, $C^*: X \rightarrow R$, $D: R \rightarrow R$.

(i) If A is nonsingular then \bar{A} is nonsingular iff

(4.2) $E = D - C^*A^{-1}B$ is nonzero.

(ii) If A is singular and $\dim N(A) = \text{codim } R(A) = 1$ then \bar{A} is nonsingular iff

(4.3) $B \notin R(A)$, $C^* \notin R(A^*)$.

Proposition 4.2 For an appropriate choice of the continuation parameter, i.e. λ respectively r , F_y in (4.1) is regular except at bifurcation points.

Proof The quantity E in (4.2) is easily computed as $E = (2-\theta)\dot{\lambda}_0 + \theta \dot{r}_0^2/\dot{\lambda}_0$ if $N = N_r$. If we accept as an appropriate choice of the continuation parameter not to pick r respectively λ to continue past a turning point in r respectively λ then we have $\theta \neq 0$ if $\dot{\lambda}_0 = 0$ respectively $\theta \neq 2$ if $\dot{r}_0 = 0$, so

that F_y is regular in regular points for $N = N_r$. In bifurcation points $G_\lambda \in R(G_u)$ so that F_y is always singular from (4.3) for $N = N_r$ and $N = N_o$. It remains to consider turning points.

If $\dot{\lambda}_0 = 0$ and $G_\lambda \notin R(G_u)$ then the last condition in (4.3) is equivalent to $(\theta \dot{r}_0 u_0 / r_0, \dot{u}_0) = \theta \dot{r}_0^2 \neq 0$. So if $\theta \neq 0$, $r_0 \neq 0$ then F_y is regular for $N = N_r$ unless $\dot{r}_0 = 0$. This situation occurs in symmetry-breaking pitchfork bifurcation points.

These points appear to be the most common type of secondary bifurcation in physical examples. Finite dimensional problems exhibiting secondary bifurcation often arise from symmetry preserving discretizations of such continuous problems. It is, therefore, important that the continuation method can handle these points. In addition to that, the proposed algorithm allows to detect these points by checking (3.14). We refer to Example 1 in [11] where two symmetry-breaking bifurcation points are computed and to the second example in 5. According to (3.15) $N = N_o$ is chosen in these points since N_r is not well defined. If the point is not a bifurcation point both conditions of (4.3) are satisfied and F_y is again regular.

The above proposition does not specify in any more detail what happens in bifurcation points. If we consider F_y^{-1} in the neighbourhood of a simple bifurcation point $y = y(s_0)$ and measure the growth of $\|F_y^{-1}\|$ as a function of $(s-s_0)^{-1}$ as in [4] then we obtain the following result along the lines of proofs in [4,5].

Proposition 4.3 If the normalization condition N is chosen according to (3.15) then on any of the branches passing through a simple bifurcation point $y_0 = (u(s_0), \lambda(s_0))$ of (2.1) for $\delta > 0$ sufficiently small

$$(4.4) \quad \|F_y^{-1}(y(s))\| = O(|s-s_0|^{-1}), \quad 0 < |s-s_0| < \delta.$$

The growth behaviour of (4.4) allows the application of the Newton-Kantorovich theorem to show convergence for the starting point obtained by the Euler predictor ([4]). For the combination with the predictor (3.2), (3.3) a different analysis would have to be done. One of the facts that complicate such an analysis considerably is the least squares solution of (3.3). We do not present such a theory here and refer to an illustration of the practical merits of the method by the numerical results in [11] and in the following section.

5. Numerical Results

We present here only a demonstration of the behaviour of the proposed continuation method. The method was developed and implemented in the program PLTMG [1] in cooperation with R. Bank. The class of second order nonlinear parameter-dependent boundary value problems solved by PLTMG includes many interesting applications.

It is of interest how the continuation method proposed above compares to standard methods as, for example, the

classical pseudo-arclength method, $\theta = 1$ in (2.3). The generalized inverse iteration ([9]) that strongly influenced the method presented here was compared to different continuation methods in [9] and to a predecessor of the present method using a two-parameter predictor and (2.5) in [10]. The latter paper also contains comparisons with a variant of the generalized inverse iteration in the neighbourhood of symmetry-breaking bifurcation points.

An extensive comparison of the present method with the pseudo-arclength method is beyond the scope of this paper. We restrict the consideration to one example exhibiting turning points and thereby illustrate the advantages of a multi-parameter predictor. The second example has a more complicated solution structure and serves to demonstrate the computation of singular points, the switching of branches at bifurcation points, the detection of symmetry-breaking bifurcation points and the switching of continuation parameters in a multi-parameter problem.

The first example is the two-parameter Bratu problem

$$(5.1) \quad \Delta u + \lambda \exp(u/(1+\epsilon u)) \text{ in } \Omega$$

on the unit square Ω with homogeneous Dirichlet boundary conditions. This problem was solved frequently (see, for example, [3,11]). For $\epsilon = .1$ the solution curve has two turning points and we present results for continuing up to and beyond the

first one. We point out that this turning point is far from being nearly angular and therefore does not cause particular difficulties for the pseudo-arclength method. For a case of a nearly angular turning point, see Example 2 in [11].

On a standard triangulation of Ω , with 25 vertices, we have discretized (5.1) by linear finite elements. The resulting finite-dimensional system was first solved by the pseudo-arclength method starting at the origin. We assumed that the step-picking procedure does not cut back the steplength α in the Euler predictor, i.e., $\alpha = \beta, \gamma = 0$ in (3.2), enough near the turning point to avoid failure of the corrector iteration. In this latter case, the steplength is cut back by .5 until the corrector converges. In the last column of Table 5.1 the number of corrector iterations is listed and an asterisk denotes failure to converge after the given number of iterations while two asterisks denote failure to yield a descent direction for the damped Newton process.

Subsequently, we have used our continuation method to continue from each of the points in Table 5.1 below the turning point directly to the point above that with $\|u\| = 2.61$ which was finally reached. The results in Table 5.2 show that at most 3 corrector steps are needed. The three predictor parameters are also given to show for this example the effect of introducing γ and the early 'detection' of the turning point

through a negative α . We note that for the coarse discretization used the first turning point in case $\epsilon = .1$ is at $(\lambda, r) = (.840, .928)$.

λ	r	$\dot{\lambda}$	\dot{r}	α	it
0	0	1	0	1	1
1	.04	.999	.042	5	2
6	.343	.995	.095	2	3
8	.646	.959	.282	.665	10*
				.332	2
8.28	.772	.847	.531	.920	10*
				.460	10**
				.230	10**
				.115	2
8.37	.848	.596	.803	1.49	10*
				.745	10*
				.373	10*
				.186	4**
				.093	2
8.40	.943	-.125	.991	2.09	4
4.91	2.61	-.894	.447		

Table 5.1 Continuation with the pseudo-arclength method for (5.1), $\epsilon = .1$

λ	$\dot{\lambda}$	α	β	γ	it
1	.999	4.40	.17E-3	64.3	3
6	.995	- .732	-.21E-2	6.58	3
8	.959	-2.96	-.51E-1	3.01	2
8.28	.847	-3.70	-.125	2.47	2
8.37	.596	-5.54	-.345	2.40	2

Table 5.2 Continuation with proposed method for each point below turning point in Table 5.1 directly to solution with norm $r = 2.61$

A suitable step-picking procedure may be able to reduce the number of failing corrector iterations for the pseudo-arclength method. But the fact remains that several small steps are necessary to overcome the turning point. The next example has a more complex solution diagram including bifurcation. It illustrates several other features of the continuation method and its implementation.

Let Ω_ϵ be a domain that is obtained by connecting two regions in the left and right half-plane situated symmetrically w.r.t. the y-axis by a channel of width ϵ such that Ω_ϵ is symmetric, too. Homogeneous Neumann boundary conditions are prescribed along $\partial\Omega_\epsilon$.

$$(5.1) \quad \frac{\partial u}{\partial t} = \Delta u + \lambda u - au^2 - u^3 \quad \text{in } \Omega_\epsilon,$$

The applications of this reaction-diffusion equation include in particular a selection-migration model from population genetics (cf. [13]). We are interested in the stationary solutions only. The bifurcation behaviour of (5.1) w.r.t. the three parameters ϵ, λ, a is given in [13]. For $a = 0$, for example, secondary bifurcation takes place from the branch bifurcating from the trivial solution at the first nonzero eigenvalue λ_ϵ of the linearization of (5.1). For $a \neq 0$, and for this case some results will be presented below, secondary symmetry-breaking bifurcation occurs from the branch of constant solutions bifurcating at the origin.

Because of symmetry and the homogeneous Neumann boundary condition, it suffices to take as Ω_ϵ a non-convex, say, L-shaped domain with vertices $(0,0)$, $(1,0)$, $(1,.5)$, $(.5,.5)$, $(.5,1)$, $(0,1)$. Thus, ϵ is fixed and the parameters a and λ are left. For $a > 0$ the following schematic bifurcation diagrams were obtained in [13] for a point value of u versus λ .




Figure 5.1a




Figure 5.1b

The elliptic version of (5.1) was discretized on a relatively coarse triangulation Ω_h with 21 vertices and linear finite elements were used to compute the approximate solution u_h . Better approximations at specified points of the solution branches may be obtained through PLTMG by a multi-grid method.

The first nontrivial eigenvalue was $\lambda_{eh} = 6.73$. In fact, for $a = 5$ no secondary bifurcation was observed. For $a = 9$ the two secondary bifurcations points in Figure 5.1b are located at $(\lambda_1, r_1) = (-8.17, 1.04)$ and $(\lambda_2, r_2) = (-14.6, 2.42)$, while the turning point is at $(\lambda_3, r_3) = (-16, 3.46)$.

We present below the output of PLTMG for the following continuation. Starting on the trivial solution in $\lambda = 0$ we switch branches three times using Method I of [6]. On the branch of constant negative solutions we continue to (λ_1, r_1) . There we switch branches once and continue on this branch of non-constant solutions to $(\lambda, r) = (-11.5, 1.6)$. Now we keep this λ fixed and switch to a as continuation parameter. From the value $a = 8$ we continue to smaller values. We expect to first come to the constant solutions and continuing further to reach a point where no further continuation is possible. The first happens between $a = 7.5$ and $a = 7.4$, as a more detailed computation shows, while the branch ends at $a = 6.78$. If instead we continue in r , then $(a, r) = (6.78, 2.94)$ is a turning point w.r.t. a as expected.

it	λ/a	$\ u\ $	$\dot{\lambda}$	\dot{r}	$\det(G_u)$	δ
Switch branches three times.						
1	0.000E+00	0.000E+00	0.100E+01	0.000E+00	-0.105E+00	-0.124E-05
0	0.000E+00	0.000E+00	0.994E+00	0.108E+00	-0.105E+00	-0.124E-05
0	0.000E+00	0.000E+00	-0.100E-01	0.000E+00	-0.105E+00	-0.124E-05
0	0.000E+00	0.000E+00	-0.994E+00	-0.108E+00	-0.105E+00	-0.124E-05
Continue to $r = 1$, then to $r = 1.5$ and check sign of determinant.						
1	-0.790E+01	0.100E+01	-0.989E+00	0.150E+00	-0.902E+02	0.435E-02
1	-0.109E+02	0.150E+01	-0.982E+00	0.188E+00	0.541E+03	-0.316E-01
Determinant changes sign; find bifurcation point by secant method on δ .						
1	-0.860E+01	0.111E+01	-0.988E+00	0.157E+00	-0.128E+03	-0.651E-02
1	-0.794E+01	0.101E+01	-0.989E+00	0.151E+00	-0.766E+02	0.370E-02
1	-0.818E+01	0.104E+01	-0.988E+00	0.153E+00	0.354E+01	-0.174E-03
Switch branches and continue to $r = 1.2$, $r = 1.6$.						
0	-0.817E+01	0.104E+01	0.484E-01	-0.639E-02	0.815E-01	-0.427E-05
2	-0.908E+01	0.116E+01	-0.980E+00	0.142E+00	-0.652E+03	0.255E-01
1	-0.931E+01	0.120E+01	-0.982E+00	0.148E+00	-0.834E+03	0.312E-01
1	-0.115E+02	0.160E+01	-0.976E+00	0.220E+00	-0.197E+04	0.633E-01
Switch continuation parameter to a . Continue to $a = 7.5$, then to $r = 3$ and check sign of determinant.						
1	0.800E+01	0.160E+01	0.829E+00	-0.419E+00	-0.197E+04	0.633E-01
3	0.750E+01	0.187E+01	0.393E+00	-0.200E+00	-0.183E+03	0.876E-02
1	0.678E+01	0.300E+01	-0.481E-01	-0.999E+00	0.162E+04	0.178E-01
Determinant changes sign; find turning point by secant method on $\det(G_u)$.						
1	0.679E+01	0.277E+01	0.146E+00	-0.989E+00	-0.218E+04	-0.469E-01
1	0.678E+01	0.294E+01	-0.398E-02	-0.100E+01	0.110E+03	0.142E-02
1	0.678E+01	0.294E+01	-0.307E-03	-0.100E+01	0.836E+01	0.109E-03
1	0.678E+01	0.294E+01	0.844E-05	-0.100E+01	-0.232E+00	-0.300E-05

Table 5.1 Continuation results for problem (5.1)

In Table 5.1, it denotes the number of (damped) Newton corrector iterations, while δ is a quantity that changes sign at simple bifurcation points. We point out that after switching onto the branch of non-constant solutions, $\dot{\lambda}$ and \dot{r} are both small, as expected, indicating a symmetry-breaking bifurcation point (cf.(3.14)).

Acknowledgement: The author would like to thank the referees for valuable comments and corrections.

References

- [1] R. E. Bank, PLTMG User's Guide, Edition 4.0, Tech. Report, Dept. Math., University of California, San Diego, 1985.
- [2] R. E. Bank and H. D. Mittelmann, Continuation and multigrid for nonlinear elliptic systems, to appear in Proceedings of the 2nd European Conference on Multigrid Methods, 1985, W. Hackbusch and U. Trottenberg, eds., Springer-Verlag, Berlin.
- [3] J. H. Bolstad and H. B. Keller, A multigrid continuation method for elliptic problems with turning points, submitted to SIAM J. Sci. Stat. Comp.
- [4] D. W. Decker and H. B. Keller, Path following near bifurcation, Comm. Pure Appl. Math. 34, 149-175 (1981).
- [5] J. Descloux, Two remarks on continuation procedures for solving some nonlinear problems, manuscript (1983).
- [6] H. B. Keller, Numerical solution of bifurcation and nonlinear eigenvalue problems, in Applications of Bifurcation Theory, P. Rabinowitz, ed., Academic Press, New York, 1977.
- [7] T. Küpper, H. D. Mittelmann and H. Weber, eds., Numerical Methods for Bifurcation Problems, ISNM 70, Birkhäuser-Verlag, Basel, 1984.
- [8] H. D. Mittelmann and H. Weber, Numerical methods for bifurcation problems - a survey and classification, in Bifurcation Problems and their Numerical Solution, H. D. Mittelmann and H. Weber, eds., ISNM 54, Birkhäuser-Verlag, Basel, 1980.
- [9] H. D. Mittelmann, A fast solver for nonlinear eigenvalue problems, in Iterative Solution of Nonlinear Systems of Equations, R. Ansorge, Th. Meis and W. Törnig, eds., Lecture Notes in Mathematics 953, Springer-Verlag, Berlin, 1982.
- [10] H. D. Mittelmann, Continuation near symmetry-breaking bifurcation points, in [5].
- [11] H. D. Mittelmann, Multi-level continuation techniques for nonlinear boundary value problems with parameter dependence, Appl. Math. Comp. (to appear).
- [12] W. C. Rheinboldt and J. V. Burkhardt, A locally parametrized continuation process, ACM Trans. Math. Software 9, 215-235 (1983).
- [13] J. M. Vegas, Bifurcations caused by perturbing the domain in an elliptic equation, J. Differential Equations 48, 189-226 (1983).

OTHER REPORTS IN THIS SERIES

Bustoz, J. and Savage, H., "Inequalities for Ultraspherical and Laguerre Polynomials," Technical Report No. 1.

Bustoz, J., "On Absolute Euler-Knopp and de la vallee Poisson Summability," Technical Report No. 2.

Busby, R.C. and Smith, H.A., "Product-Convolution Operators and Mixed-Norm Spaces," Technical Report No. 3.

Kupperman, R. H. and Smith, H. A., "The Effects of MIRV on Force Size in Mutual Deterrence," Technical Report No. 4.

Aickin, Mikel, "Linear Analysis on Contingency Tables," Technical Report No. 5.

Grace, E. E., "Refinable maps on graphs are near homeomorphisms," Technical Report No. 6.

Kuiper, H. S. and Shew, S. H., "Strong Solutions for Infinite-Dimensional Riccati Equations Arising in Transport Theory," Technical Report No. 7.

Atalla, Robert E., "Generalized Almost-Convergence vs. Matrix Summability," Technical Report No. 8.

Anderson, B. A. and Gross, K. B., "A Partial Starter construction," Technical Report No. 9.

McDonald, John D., "Approximation by quotients of rational inner functions," Technical Report No. 10.

Anderson, B. A., Gross, K. B., and Leonard, P. A., "Some Howell Designs of Prime Size," Technical Report No. 11.

Wang, Cecilia Y., "Stiffness of Harmonic Functions," Technical Report No. 12.

Wang, Alan P., "Dissipative Properties of A Nonlinear Differential Operators Equation," Technical Report No. 13.

Aickin, Mikel, "Existence of MLEs for Discrete Linear Exponential Models," Technical Report No. 14.

Wang, Cecilia Y., "A biharmonic kernel function," Technical Report No. 15.

Chung, Lung Ock, Sario, Leo, and Wang, Cecilia, "Harmonic L^p functions on the Poincare D-ball," Technical Report No. 16.

Talaga, Paul C., "Invariant Sets, Existence Theorems, and the Hukuhara-Kneser Property for Semilinear Parabolic Systems with Neumann Type boundary Condition," Technical Report No. 17.

Baumeister, Richard, "Applications of Clebsch Potentials to Variational Principles in the Theory of Physical Fields," Technical Report No. 18.

Kuiper, Hendrik J., "Invariant Sets for Nonlinear Elliptic and Parabolic Systems," Technical Report No. 19.

McMahon, Douglas, "Relativized weak mixing of uncountable order," Technical Report No. 20.

Bustoz, J. and Savage, H., "Inequalities for ultraspherical and Laguerre polynomials II," Technical Report No. 21.

Schwartz, David P., "Time-sliced Multi-tasking in a two-job Environment" (Communicated by Professor Alan Feldstein, Dept. of Mathematics, ASU, Tempe, AZ) Technical Report No. 22.

Waltman, Paul, Hubbell, Stephen P. and Hsu, Sze-Bi, "Theoretical and Experimental Investigation of Microbial Competition in Continuous Culture," Technical Report No. 23.

Anderson, B. A., "Hyperovals and Howell Designs," Technical Report No. 24.

Anderson, B. A. and Leonard, P. A., "Sequencings and Howell Designs," Technical Report No. 25.

Anderson, B. A., "A Doubling Construction for Certain Howell Designs," Technical Report No. 26.

Feldstein, A. and Schwartz, D., "A Beginner's Guide to Tape Sabotage," Technical Report No. 27.

Leonard, Philip A. and Williams, Kenneth S., "The Spinor Genera for a Certain Form Discriminant," Technical Report No. 28.

Anderson, B. A., "Some Howell Designs of Side 6P," Technical Report No. 29.

McDonald, John N., "On Some Zero Configurations Associated with the van der Waerden Conjecture," Technical Report No. 30.

Kelly, John B., "Difference Systems, Graph Designs, and Coloring Problems," Technical Report No. 31.

Bustoz, J., "Two parameter Turan inequalities for Ultraspherical and Laguerre polynomials," Technical Report No. 32.

Ismail, Mourad E.H., "An Asymptotic Result for a Learning Model," Technical Report No. 33.

Feldstein, Alan and Neves, Kenneth W., "High Order Methods for State Dependent Delay Differential Equations With Nonsmooth Solutions," Technical Report No. 34.

Anderson, B. A. and Gross, K. B., "Asymptotic Multiplication of Howell Designs by 3," Technical Report No. 35.

Anderson, B. A., "A Note on Howell Designs," Technical Report No. 36.

Anderson, B. A. and Leonard, P. A., "Some Howell Designs of Prime Side II," Technical Report No. 37.

McDonald, J. N., "Closed Orbits of Convex Sets of Operators on the Disk Algebra," Technical Report No. 38.

Young, Dennis, "The Linear Nearest Neighbor Statistic," Technical Report No. 39.

Ismail, Hourad E.H., "The Basic Bessel Functions and Polynomials," Technical Report No. 40.

Aickin, Mikel, "DIGRAPH : A program for Testing Multiple Hypotheses," June 1980, Technical Report No. 41.

Salamon, Peter, Bjarne Andresen, Paul D. Gait, and R. Stephen Berry, "The Significance of Weinhold's Length", June 1980, Technical Report No. 42.

Salamon, Peter, and A. Nitzan, "Finite Time Optimizations of A Newton's Law Carnot Cycle," June 1980, Technical Report No. 43.

Salamon, Peter, Yehuda B. Band, and Oded Kafri, "Finite Time Thermodynamics: Optimal Expansion of A Heated Working Fluid," June 1980, Technical Report No. 44.

Salamon, Peter, Y. B. Band and O. Kafri, "Maximum Power From A Cycling Working Fluid," June 1980, Technical Report No. 45.

Salamon, Peter, Yehuda B. Band, and Oded Kafri, "Maximum Work Production from a Heated Gas in a Cylinder Equipped with a Piston," June 1980, Technical Report No. 46.

Jacobowitz, Ronald and Driscoll, Michael F., "Estimability Testability, and Parametrization in Fixed-Effects Linear Models," July 1980, Technical Report No. 47.

Dudley, David, and Wang, Alan, P., "Operator Theory on the WKE Method and Bremmer Series," July 1980, Technical Report No. 48.

Ihrig, Edwin C. and Ismail, Mourad E.H., "A q-Umbral Calculus*", August 1980, Technical Report No. 49.

Smith, Hal, "The interaction of steady state and Hopf bifurcations in a two predator - one prey competition model," November 1980, Technical Report No. 50.

Grace, Edward E., "Local Properties in II_2 Spaces," December 1980, Technical Report No. 51.

Grace, Edward E., "Cuts points in totally non-semi-locally-connected II_2 continua," December 1980, Technical Report No. 52.

Bustoz, Joaquin and Ismail, Mourad E.H., "The associated ultraspherical polynomials and their q-analogues," February 1981, Technical Report No. 53.

Aickin, Mikel, "HLA User's Handbook," February 1981, Technical Report No. 54.

Driscoll, Michael F. and Young, Dennis L., "Some Tests for Assessing Randomness in One or More Dimensions," April 1981, Technical Report No. 55.

Buzano, Ernesto, Golubitsky, Martin, "Bifurcation Involving the Hexagonal Lattices," August 1981, Technical Report No. 56.

Anderson, B. A., "Hyperplanes and Balanced Howell Rotations," August 1981, Technical Report No. 57.

Ismail, M.E.H. and Wilson, James A., "Asymptotic and Generating Relations for the q -Jacobi and q, ϕ_3 Polynomials," Technical Report No. 58., August 1981.

Ismail, M.E.H. and Miller Kenneth S., "An Infinitely Divisible Distribution Involving Modified Bessel Functions", Technical Report No. 59.

Keyfitz, Barbara and Kuiper, Hendrik, "Bifurcation Resulting from Changes in Domain in a Reaction-Diffusion Equation*," Technical Report No. 60.

Herrero, Domingo A., "Compact Perturbations of Nest Algebras, Index Obstructions and A Problem of Arveson," Technical Report No. 61.

Keyfitz, Barbara and Kuiper, Hendrik, "Bifurcation Resulting from Changes in Domain in a Reaction-Diffusion Equation*," Technical Report No. 60.

Bustoz, J. and Ismail, Mourad E.H., "Turan Type Inequalities for Ultraspherical, Continuous q -Ultraspherical, and Hermite Polynomials," Technical Report No. 63.

Wang, Alan P., "Solutions of a Dissipative Diffusion Equation in a Hilbert Space," Technical Report No. 64, January, 1982.

Smith, Hal L., "Subharmonic Bifurcation in an S-I-R Epidemic Model", Technical Report No. 65, February 1982.

Goldstein, M. and Herrero, D., "Infinite Blaschke Products Which Almost Omit a Dense Set of Values", Technical Report No. 66, March, 1982.

Goldstein, M. and McDonald, J. N., "An extremal problem for non-negative trigonometric polynomials", Technical Report No. 67, March, 1982.

Herrero, Domingo, A., "Three Results on approximation of Hilbert Space Operators", "Economical Compact Perturbations I: Erasing Normal Eigenvalues", "Compact Perturbations of Continuous Nest Algebras", Technical Report No. 68, March, 1982.

Kuiper, Hendrik J., "Generalized Operator Riccati Equations", Technical Report No. 69, April, 1982.

Kelly, John B., "Facets of the Hamming Cone", Technical Report No. 70, May, 1982.

Herrero, D., "On Limits of Nilpotents in LP -spaces and Interpolation of Spectra", Technical Report NO. 71, July, 1982.

Herrero, D., "Two Results on Bilateral Weighted Shifts", Technical Report No. 72, September, 1982.

Herrero, D. and McDonald, J., "On Multicyclic Operators and the Vasjuninnikol'skii Discotheca", Technical Report No. 73, September, 1982.

Herrero, D., "On Multicyclic Operators. II: Two Extensions of the Notion of Quasitriangularity", Technical Report No. 74, November 1982.

McDonald, John N., "Examples of RP -measures", Technical Report No. 75, May 1983.

Smith, Hal. L., "Normal Forms for Periodic Systems", Technical Report No. 76, July 1983.

Ismail, Mourad E.H., "On Sieved Orthogonal Polynomials", Technical Report No. 77, August 1983.

Ismail, Mourad E.H. and Askey, Richard A., "Recurrence Relations, Continued Fractions and Orthogonal Polynomials", Technical Report No. 78, September, 1983.

Mittelmann, H.D., "Continuation Near Symmetry-Breaking Bifurcation Points", Technical Report No.79, October 1983.

Miel, George, "Perturbed Projection Methods for Split Equations of the First Kind", Technical Report No. 80, December, 1983.

Miel, George, "On the Galerkin and Collocation Methods for a Cauchy Singular Integral Equation", Technical Report No. 81, December, 1983.

McDonald, John N., "Unimodular Approximation in Function Algebras", Technical Report No. 82, March 1984.

Ringhofer, Christian A., "A Second Order Difference Scheme for Transient Semiconductor Device Simulation", Technical Report No. 83, January, 1985.

Mittelmann, Hans D., "A Free Boundary Problem and Stability for the Nonlinear Beam", Technical Report No. 84, March 1985.

Mittelmann, Hans D., "Multi-Level Continuation Techniques for Nonlinear Boundary Value Problems with Parameter Dependence", Technical Report No. 85, March 1985.

Kuiper, Hendrik J., "On Spurious Numerical Solutions for Nonlinear Eigenvalue Problems," Technical Report No. 86, July 1985.

Mittelmann, Hans D., "Continuation and Multi-grid for Nonlinear Elliptic Systems," Technical Report No. 87, September 1985.

Mittelmann, Hans D., "A Pseudo-Arclength Continuation Method for Nonlinear Eigenvalue Problems," Technical Report No. 88, October 1985.

END

12-86

DTIC